arXiv:1006.3025v2 [math.NT] 27 Jun 2010

SOME CONGRUENCES FOR TRINOMIAL COEFFICIENTS

HUI-QIN CAO AND HAO PAN

Abstract. We prove several congruences for trinomial coefficients.

1. Introduction

In [3], Pan and Sun proved the following congruence on the sums of binomial coefficients:

$$\sum_{k=0}^{p-1}\binom{2k}{k+d}\equiv\left(\frac{p-d}{3}\right)\pmod p, \tag{1.1}$$

where $p>3$ is a prime, $0\leqslant d\leqslant p-1$ and $\left(\frac{-}{}\right)$ is the Legendre symbol. They also proved that for prime $p>3$ and integer $0\leqslant d\leqslant p-1$,

$$\sum_{k=1}^{p-1}\frac{1}{k}\binom{2k}{k+d}\equiv\begin{cases}d^{-1}(-1+2(-1)^d+3[3\mid p-d])\pmod p, & \text{if } 1\leqslant d\leqslant p,\\ 0\pmod p, & \text{if } d=0,\end{cases} \tag{1.2}$$

where $[A]=1$ or 0 according to whether the assertion A holds. Subsequently, Sun and Tauraso [6] extended (1.1) and showed that

$$\sum_{k=0}^{p-1}\binom{2k}{k+d}\equiv\left(\frac{p-d}{3}\right)+2pS_d\pmod{p^2}, \tag{1.3}$$

where

$$S_d=\sum_{0<k<d}\frac{(-1)^{k-1}}{k}\left(\frac{d-k}{3}\right).$$

On the other hand, trinomial coefficients $\binom{n}{k}_2$ are given by

$$(1+x+x^{-1})^n=\sum_{k=-n}^{n}\binom{n}{k}_2x^k.$$

As G. E. Andrews mentioned [1], trinomial coefficients had been investigated by Euler. And Andrews and R. J. Baxter [2] found the q-analogues of trinomial coefficients play an important rule in the hard hexagon model. However, there is a similar congruence for trinomial coefficients:

$$\sum_{k=0}^{p-1}\binom{k}{d}_2\equiv\begin{cases}(-1)^{\frac{p+d-1}{2}}\pmod p & \text{if } d \text{ is even},\\ 0\pmod p & \text{if } d \text{ is odd},\end{cases} \tag{1.4}$$

where $p>3$ is prime and $0\le d<p$. In fact, we shall prove the following stronger result.

Theorem 1.1. *Let $p > 3$ be a prime and let d be an integer with $0 \le d \le p-1$. If d is odd, then*

$$\frac{1}{p}\sum_{k=0}^{p-1}\binom{k}{d}_2 \equiv \frac{(-1)^{\frac{d+1}{2}}}{2}\Bigg(\sum_{k=1}^{(d-1)/2}\frac{(-1)^k}{k} - 3\sum_{\substack{1\le k\le (d-1)/2\\ 3\mid k+p}}\frac{(-1)^k}{k}\Bigg) \pmod{p}. \tag{1.5}$$

And if d is even, then

$$\begin{aligned}&\frac{1}{p}\Bigg((-1)^{\frac{d}{2}}\sum_{k=0}^{p-1}\binom{k}{d}_2 - (-1)^{\frac{p-1}{2}}\Bigg)\\ \equiv &-3\left(\frac{-2}{p}\right)\frac{\mathcal{S}_{(p-(\frac{3}{p}))/2}}{p} - 2\sum_{0\le j<d/2}\frac{(-1)^j}{2j+1} + 3\sum_{\substack{0\le j<d/2\\ 3\nmid p-2j-1}}\frac{(-1)^j}{2j+1} \pmod{p},\end{aligned} \tag{1.6}$$

where the recurrence sequence $\{\mathcal{S}_n\}$ is defined as

$$\mathcal{S}_0 = 0, \quad \mathcal{S}_1 = 1, \quad \mathcal{S}_{n+1} = 4\mathcal{S}_n - \mathcal{S}_{n-1} \text{ for } n \ge 1.$$

We also have a congruence for the alternating sums of trinomial coefficients.

Theorem 1.2. *Let $p > 3$ be a prime and let d be an integer with $0 \le d \le p-1$. We have*

$$\sum_{k=0}^{p-1}(-1)^k\binom{k}{d}_2 \equiv (-1)^{d-1}d + (-1)^d p\sum_{k=d+1}^{\lfloor (p+d-1)/2\rfloor}\frac{1}{k}\binom{2k-d}{k+1} \pmod{p^2}. \tag{1.7}$$

Unfornately, in general, it is not easy to compute

$$\sum_{k=d+1}^{\lfloor (p+d-1)/2\rfloor}\frac{1}{k}\binom{2k-d}{k+1} \pmod{p}.$$

However, when $d = 0, 1, 2$, we may get

Theorem 1.3. *For prime $p > 3$ we have*

$$\sum_{k=0}^{p-1}(-1)^k\binom{k}{0}_2 \equiv p\frac{3\left(\frac{p}{3}\right)-1}{2} \pmod{p^2}, \tag{1.8}$$

$$\sum_{k=0}^{p-1}(-1)^k\binom{k}{1}_2 \equiv 1 - p\frac{3\left(\frac{p}{3}\right)-1}{2} \pmod{p^2} \tag{1.9}$$

and

$$\sum_{k=0}^{p-1}(-1)^k\binom{k}{2}_2 \equiv -2 + 3p\left(\frac{p}{3}\right) \pmod{p^2}. \tag{1.10}$$

Quiet recently, Z.-W. Sun told us that with help of the software *Mathematica*, he found a similar congruence for the sum

$$\sum_{k=0}^{p-1}\frac{T_k}{3^k}.$$

Here we shall give a proof of Sun's congruence.

Theorem 1.4. *Let $p > 3$ be a prime. Then*

$$\sum_{k=0}^{p-1} \frac{T_k}{3^k} \equiv \begin{cases} p \pmod{p^2} & \text{if } p \equiv 1 \pmod 3, \\ 0 \pmod{p^2} & \text{if } p \equiv 2 \pmod 3. \end{cases}$$

The proofs of the above theorems will be given in the next sections.

2. Proof of Theorem 1.1

Lemma 2.1. *Suppose that $n > d \geq 0$. Then*

$$\sum_{k=0}^{n-1} \binom{k}{d}_2 = \sum_{\substack{k=1 \\ k+d \equiv 1 \pmod 2}}^{n} \binom{n}{k} \binom{k-1}{\frac{k+d-1}{2}}. \tag{2.1}$$

Proof. Let $[x^d]P(x)$ denote the coefficient of x^d in the expansion of the polynomial $P(x)$. Since

$$\sum_{k=0}^{n-1} (1+x+x^{-1})^k = \frac{(1+x+x^{-1})^n - 1}{x+x^{-1}} = \frac{1}{x^{n-1}} \cdot \frac{(1+x+x^2)^n - x^n}{1+x^2},$$

we have

$$\begin{aligned} \sum_{k=0}^{n-1} \binom{k}{d}_2 =& \sum_{k=0}^{n-1} [x^d](1+x+x^{-1})^k = [x^d] \frac{1}{x^{n-1}} \cdot \frac{(1+x+x^2)^n - x^n}{1+x^2} \\ =& [x^{n+d-1}] \frac{(1+x+x^2)^n - x^n}{1+x^2} = [x^{n+d-1}] \sum_{k=1}^{n} \binom{n}{k} (1+x^2)^{k-1} x^{n-k} \\ =& \sum_{k=1}^{n} \binom{n}{k} [x^{k+d-1}](1+x^2)^{k-1} = \sum_{\substack{k=1 \\ k+d \equiv 1 \pmod 2}}^{n} \binom{n}{k} \binom{k-1}{\frac{k+d-1}{2}}. \end{aligned}$$

□

Noting that

$$\binom{p}{k} \equiv 0 \pmod p \qquad \text{and} \qquad \binom{p-1}{k} \equiv (-1)^k \pmod p$$

for $0 < k < p$, (1.4) immediately follows from (2.1).

Lemma 2.2. *Suppose that $p > 3$ is prime and $0 \leq m < (p-1)/2$. Then*

$$\sum_{j=1}^{(p-1)/2} \frac{1}{j} \binom{2j}{j+m} \equiv \frac{1}{m}(1 - 3[3 \mid p+m]) \pmod p.$$

Proof. Let

$$a_m = \sum_{j=1}^{(p-1)/2} \frac{1}{j} \binom{2j}{j+m}.$$

Define the polynomials $\{v_n(x)\}_{n\in\mathbb{N}}$ by

$$v_0(x)=2, v_1(x)=x, \text{ and } v_{n+1}(x)=xv_n(x)-v_{n-1}(x), \quad n=1,2,\dots.$$

Applying Theorem 2.1 in [5], we have

$$\begin{aligned}m\sum_{j=1}^{\frac{p-1}{2}}\frac{1}{j}\binom{2j}{j+m}-v_m(-1)=&-\sum_{j=0}^{\frac{p-1}{2}+m}\binom{p+1}{j}v_{\frac{p+1}{2}+m-j}(-1)-2\binom{p}{\frac{p-1}{2}+m}\\ \equiv&-v_{\frac{p+1}{2}+m}(-1)-(p+1)v_{\frac{p-1}{2}+m}(-1)\\ \equiv&-v_{\frac{p+1}{2}+m}(-1)-v_{\frac{p-1}{2}+m}(-1)=v_{\frac{p+1}{2}+m+1}(-1) \pmod p.\end{aligned}$$

Since $v_n(-1)=3[3\mid n]-1$ for all $n\in\mathbb{N}$, we have

$$\begin{aligned}a_m=\sum_{j=1}^{\frac{p-1}{2}}\frac{1}{j}\binom{2j}{j+m}\equiv&\frac{1}{m}(v_m(-1)+v_{\frac{p+3}{2}+m}(-1))\\ =&\frac{1}{m}(3[3\mid m]+3[3\mid \frac{p+3}{2}+m]-2)\\ =&\frac{1}{m}(3[3\mid m]+3[3\mid p-m]-2)\\ =&\frac{1}{m}(1-3[3\mid p+m]) \pmod p.\end{aligned}$$

□

Lemma 2.3. *Suppose that $p>3$ is prime and $0\le m<(p-1)/2$. Then*

$$\sum_{j=1}^{(p-3)/2}\frac{(-1)^j}{2j+1}\equiv\left(\frac{-1}{p}\right)\frac{2^{p-1}-1}{2p} \pmod p.$$

Proof. Clearly,

$$\sum_{j=1}^{(p-3)/2}\frac{(-1)^j}{2j+1}\equiv\sum_{j=1}^{(p-3)/2}\frac{(-1)^j}{2j+1}\binom{p-1}{2i}=\frac{1}{p}\sum_{j=1}^{(p-3)/2}(-1)^j\binom{p}{2j+1} \pmod p.$$

And

$$\sum_{j=1}^{(p-3)/2}(-1)^j\binom{p}{2j+1}=(-1)^{\lfloor\frac{p-1}{4}\rfloor}2^{\frac{p-1}{2}}-(-1)^{\frac{p-1}{2}}=\left(\frac{-2}{p}\right)\left(2^{\frac{p-1}{2}}-\left(\frac{2}{p}\right)\right).$$

Finally,

$$\begin{aligned}2^{p-1}-1=&\left(2^{\frac{p-1}{2}}-\left(\frac{2}{p}\right)\right)\left(2^{\frac{p-1}{2}}+\left(\frac{2}{p}\right)\right)\\ =&\left(2^{\frac{p-1}{2}}-\left(\frac{2}{p}\right)\right)^2+2\left(\frac{2}{p}\right)\left(2^{\frac{p-1}{2}}-\left(\frac{2}{p}\right)\right)\equiv2\left(\frac{2}{p}\right)\left(2^{\frac{p-1}{2}}-\left(\frac{2}{p}\right)\right) \pmod{p^2}.\end{aligned}$$

□

Lemma 2.4. *Suppose that $p > 3$ is prime. Then*

$$\sum_{\substack{0\le j\le (p-3)/2\\ 3\nmid p-2j-1}} \frac{(-1)^j}{2j+1} \equiv \left(\frac{-1}{p}\right)\frac{2^{p-1}-1}{3p} - \left(\frac{-2}{p}\right)\frac{\mathcal{S}_{(p-(\frac{3}{p}))/2}}{p} \pmod p.$$

Proof. This is an immediate consequence of [4, Corollary 3.3]. □

Now we are ready to prove (1.5) and (1.6).

Proof of (1.5). Suppose that $d = 2m+1$. Let

$$S_m = \frac{1}{p}\sum_{k=0}^{p-1}\binom{k}{2m+1}_2.$$

According to Lemma 2.1,

$$\begin{aligned} S_m = \frac{1}{p}\sum_{\substack{k=1\\ 2\mid k}}^{p}\binom{p}{k}\binom{k-1}{\frac{k+2m}{2}} &= \sum_{j=1}^{(p-1)/2}\frac{1}{2j}\binom{p-1}{2j-1}\binom{2j-1}{j+m}\\ &\equiv -\frac{1}{2}\sum_{j=1}^{(p-1)/2}\frac{1}{j}\binom{2j-1}{j+m} \pmod p. \end{aligned}$$

By (1.2),

$$\sum_{j=1}^{p-1}\frac{1}{j}\binom{2j}{j} \equiv 0 \pmod p.$$

As

$$\binom{2j}{j} \equiv 0 \pmod p$$

for $(p-1)/2 < j < p$, we obtain that

$$\begin{aligned} S_0 \equiv -\frac{1}{2}\sum_{j=1}^{(p-1)/2}\frac{1}{j}\binom{2j-1}{j} &= -\frac{1}{4}\sum_{j=1}^{(p-1)/2}\frac{1}{j}\binom{2j}{j}\\ \equiv -\frac{1}{4}\sum_{j=1}^{p-1}\frac{1}{j}\binom{2j}{j} &\equiv 0 \pmod p. \end{aligned}$$

For $1\le m\le (p-3)/2$,

$$\begin{aligned}S_m\equiv&-\frac{1}{2}\sum_{j=1}^{(p-1)/2}\frac{1}{j}\left(\binom{2j}{j+m}-\binom{2j-1}{j+m-1}\right)\\=&-\frac{1}{2}\sum_{j=1}^{(p-1)/2}\frac{1}{j}\binom{2j}{j+m}+\frac{1}{2}\sum_{j=1}^{(p-1)/2}\frac{1}{j}\binom{2j-1}{j+m-1}\\\equiv&-\frac{1}{2}\sum_{j=1}^{(p-1)/2}\frac{1}{j}\binom{2j}{j+m}-S_{m-1}\pmod p.\end{aligned}$$

Then

$$S_m\equiv -S_{m-1}-\frac{1}{2}\sum_{j=1}^{(p-1)/2}\frac{1}{j}\binom{2j}{j+m}\pmod p,$$

and therefore

$$\begin{aligned}S_m\equiv&(-1)^mS_0+\frac{(-1)^{m+1}}{2}\sum_{k=1}^{m}(-1)^k\sum_{j=1}^{(p-1)/2}\frac{1}{j}\binom{2j}{j+k}\\\equiv&\frac{(-1)^{m+1}}{2}\sum_{k=1}^{m}(-1)^k\sum_{j=1}^{(p-1)/2}\frac{1}{j}\binom{2j}{j+k}\\\equiv&\sum_{k=1}^{m}\frac{(-1)^k}{k}(1-3[3\mid p+k])=\sum_{k=1}^{m}\frac{(-1)^k}{k}-3\sum_{\substack{k=1\\3\mid k+p}}^{m}\frac{(-1)^k}{k}\pmod p.\end{aligned}$$

This concludes the proof. □

Proof of (1.6). Suppose that $d=2m$. Note that

$$\binom{k+1}{2m+1}_2=\binom{k}{2m}_2+\binom{k}{2m+1}_2+\binom{k}{2m+2}_2.$$

Hence

$$\sum_{k=0}^{p-1}\binom{k}{2m}_2+\sum_{k=0}^{p-1}\binom{k}{2m+2}_2=\sum_{k=0}^{p-1}\binom{k+1}{2m+1}_2-\sum_{k=0}^{p-1}\binom{k}{2m+1}_2=\binom{p}{2m+1}_2.$$

Let

$$S_m=\sum_{k=0}^{p-1}\binom{k}{2m}_2.$$

Then for every $j\ge 1$,

$$S_m=(-1)^jS_{m+j}+\sum_{i=0}^{j-1}(-1)^i\binom{p}{2m+2i+1}_2$$

In particular, when $j=(p-1)/2-m$, we have

$$S_m=(-1)^{(p-1)/2-m}+\sum_{i=0}^{(p-3)/2-m}(-1)^i\binom{p}{2m+2i+1}_2.$$

Now

$$\begin{aligned}\frac{1}{p}\binom{p}{2m+2i+1}_2&=\frac{1}{p}\sum_{k=0}^{p}(-1)^k\binom{p}{k}\binom{2(p-k)}{p-k-(2m+2i+1)}\\ =&\frac{2}{p-2m-2i-1}\binom{2p-1}{p-2m-2i-2}+\sum_{k=1}^{p-1}(-1)^k\frac{1}{k}\binom{p-1}{k-1}\binom{2(p-k)}{p-k-2m-2i-1}\\ \equiv&\frac{2}{2m+2i+1}+\sum_{k=1}^{p-1}\frac{1}{p-k}\binom{2(p-k)}{p-k-2m-2i-1}\pmod p.\end{aligned}$$

Theorefore

$$\begin{aligned}&\frac{S_m-(-1)^{(p-1)/2-m}}{p}\\ \equiv&\sum_{i=0}^{(p-3)/2-m}(-1)^i\bigg(\frac{2}{2m+2i+1}+\sum_{k=1}^{p-1}\frac{1}{k}\binom{2k}{k-2m-2i-1}\bigg)\\ \equiv&\sum_{i=0}^{(p-3)/2-m}(-1)^i\bigg(\frac{2}{2m+2i+1}+\sum_{k=1}^{p-1}\frac{1}{k}\binom{2k}{k+2m+2i+1}\bigg)\\ \equiv&\sum_{i=0}^{(p-3)/2-m}(-1)^i\bigg(\frac{2}{2m+2i+1}+\frac{3}{2m+2i+1}([3\nmid p-(2m+2i+1)]-1)\bigg)\\ =&2\sum_{i=0}^{(p-3)/2-m}\frac{(-1)^i}{2m+2i+1}-3\sum_{\substack{0\le i\le(p-3)/2-m\\ 3\nmid p-(2m+2i+1)}}\frac{(-1)^i}{2m+2i+1}\pmod p.\end{aligned}$$

By Lemma 2.3,

$$\sum_{i=0}^{(p-3)/2-m}\frac{(-1)^{m+i}}{2m+2i+1}\equiv\left(\frac{-1}{p}\right)\frac{2^{p-1}-1}{2p}-\sum_{j=0}^{m-1}\frac{(-1)^j}{2j+1}\pmod p.$$

And by Lemma 2.4,

$$\begin{aligned}&\sum_{\substack{0\le i\le(p-3)/2-m\\ 3\nmid p-(2m+2i+1)}}\frac{(-1)^{m+i}}{2m+2i+1}\\ \equiv&\left(\frac{-1}{p}\right)\frac{2^{p-1}-1}{3p}-\left(\frac{-2}{p}\right)\frac{\mathcal{S}_{(p-(\frac{3}{p}))/2}}{p}-\sum_{\substack{0\le j\le m-1\\ 3\nmid p+2j+1}}\frac{(-1)^j}{2j+1}\pmod p.\end{aligned}$$

We are done. □

3. Proofs of Theorems 1.2 - 1.4

Proof of Theorem 1.2. Note that

$$\sum_{k=0}^{p-1}(-1)^k\binom{k}{p-1}_2=(-1)^{p-1}\binom{p-1}{p-1}_2=1$$

and

$$\sum_{k=0}^{p-1}(-1)^k\binom{k}{p-2}_2=(-1)^{p-2}\binom{p-2}{p-2}_2+(-1)^{p-1}\binom{p-1}{p-2}_2=-1+p-1=p-2.$$

So we may assume that $d<p-2$. Since

$$\sum_{k=0}^{p-1}(-1)^k(1+x+x^{-1})^k=\frac{1+(1+x+x^{-1})^p}{1+(1+x+x^{-1})}=\frac{x^p+(1+x+x^2)^p}{x^{p-1}(1+x)^2},$$

we have

$$\begin{aligned}\sum_{k=0}^{p-1}(-1)^k\binom{k}{d}_2=&\sum_{k=0}^{p-1}(-1)^k[x^d](1+x+x^{-1})^k=[x^d]\sum_{k=0}^{p-1}(-1)^k(1+x+x^{-1})^k\\=&[x^{p+d-1}]\frac{x^p+(1+x+x^2)^p}{(1+x)^2}\\=&[x^{d-1}]\frac{1}{(1+x)^2}+[x^{p+d-1}]\sum_{k=0}^{p}\binom{p}{k}x^{2k}(1+x)^{p-k-2}\\=&(-1)^{d-1}d+\sum_{k=d+1}^{\lfloor(p+d-1)/2\rfloor}\binom{p}{k}\binom{p-k-2}{p+d-1-2k}\\=&(-1)^{d-1}d+p\sum_{k=d+1}^{\lfloor(p+d-1)/2\rfloor}\frac{1}{k}\binom{p-1}{k-1}\binom{p-k-2}{k-d-1}.\end{aligned}$$

Observe that

$$\binom{p-1}{k-1}\equiv(-1)^{k-1}\pmod p$$

and

$$\begin{aligned}\binom{p-k-2}{k-d-1}=&\frac{(p-k-2)(p-k-3)\cdots(p-2k+d)}{(k-d-1)!}\\\equiv&(-1)^{k-d-1}\frac{(k+2)(k+3)\cdots(2k-d)}{(k-d-1)!}\\=&(-1)^{k-d-1}\binom{2k-d}{k-d-1}=(-1)^{k-d-1}\binom{2k-d}{k+1}\pmod p.\end{aligned}$$

So (1.7) is valid. □

Proof of Theorem 1.3. Let $C_n=\frac{1}{n+1}\binom{2n}{n}$ be the Catalan number. Applying (1.7) with $d=0$, d=1 and $d=2$ respectively, we get

$$\sum_{k=0}^{p-1}(-1)^k\binom{k}{0}_2\equiv p\sum_{k=1}^{(p-1)/2}\frac{1}{k}\binom{2k}{k+1}=p\sum_{k=1}^{(p-1)/2}C_k \pmod{p^2},$$

$$\sum_{k=0}^{p-1}(-1)^k\binom{k}{1}_2\equiv 1-p\sum_{k=2}^{\lfloor p/2\rfloor}\frac{1}{k}\binom{2k-1}{k+1}=1-\frac{p}{2}\sum_{k=2}^{(p-1)/2}\frac{k-1}{k}C_k \pmod{p^2}$$

and

$$\begin{aligned}\sum_{k=0}^{p-1}(-1)^k\binom{k}{2}_2&\equiv -2+p\sum_{k=3}^{(p+1)/2}\frac{1}{k}\binom{2k-2}{k+1}=-2+p\sum_{k=3}^{(p+1)/2}C_{k-1}\frac{(k-2)(k-1)}{k(k+1)}\\&=-2+p\Big(\sum_{k=3}^{(p+1)/2}C_{k-1}-\sum_{k=3}^{(p+1)/2}C_{k-1}\frac{4k-2}{k(k+1)}\Big)\\&=-2+p\Big(\sum_{k=2}^{(p-1)/2}C_k-\sum_{k=3}^{(p+1)/2}\frac{C_k}{k}\Big) \pmod{p^2}.\end{aligned}$$

In [3], Pan and Sun have proved that

$$\sum_{k=0}^{p-1}C_k\equiv\frac{3(\frac{p}{3})-1}{2} \pmod p \quad\text{and}\quad \sum_{k=1}^{p-1}\frac{C_k}{k}\equiv\frac{3}{2}\Big(1-\Big(\frac{p}{3}\Big)\Big) \pmod p.$$

Clearly,

$$C_0=C_1=1,\quad C_2=2,\quad C_{p-1}=\frac{1}{p}\binom{2p-2}{p-1}\equiv -1 \pmod p$$

and

$$C_k\equiv 0 \pmod p\quad\text{for}\quad (p-1)/2<k<p-1.$$

Therefore

$$\sum_{k=1}^{(p-1)/2}C_k\equiv\sum_{k=0}^{p-1}C_k\equiv\frac{3(\frac{p}{3})-1}{2} \pmod p$$

and

$$\sum_{k=1}^{(p-1)/2}\frac{C_k}{k}\equiv\frac{1-3(\frac{p}{3})}{2} \pmod p.$$

Hence

$$\sum_{k=2}^{(p-1)/2}\frac{k-1}{k}C_k=\sum_{k=1}^{(p-1)/2}C_k-\sum_{k=1}^{(p-1)/2}\frac{C_k}{k}\equiv 3\Big(\frac{p}{3}\Big)-1 \pmod p$$

and

$$\sum_{k=2}^{(p-1)/2} C_k - \sum_{k=3}^{(p+1)/2} \frac{C_k}{k} \equiv \frac{3\binom{p}{3}-1}{2} - 1 - \frac{1-3\binom{p}{3}}{2} + 2 = 3\left(\frac{p}{3}\right) \pmod{p}.$$

This yields (1.8), (1.9) and (1.10). We are done. □

Proof of Theorem 1.4. Clearly

$$\begin{aligned}&\sum_{k=0}^{p-1}\left(\frac{(1+x+x^{-1})}{3}\right)^k = \frac{(1+x+x^{-1})^p/3^p-1}{(1+x+x^{-1})/3-1}\\ =&\frac{1}{3^{p-1}x^{p-1}}\cdot\frac{(1+x+x^2)^p-3^px^p}{(1-x)^2}\\ =&\frac{1}{3^{p-1}x^{p-1}}\cdot\frac{(1-x^3)^p-(3x(1-x))^p}{(1-x)^{p+2}}.\end{aligned}$$

Then

$$\begin{aligned}&\sum_{k=0}^{p-1}\frac{T_k}{3^k} = [x^0]\sum_{k=0}^{p-1}\left(\frac{(1+x+x^{-1})}{3}\right)^k\\ =&\frac{1}{3^{p-1}}[x^{p-1}]\frac{(1-x^3)^p-(3x(1-x))^p}{(1-x)^{p+2}}\\ =&\frac{1}{3^{p-1}}[x^{p-1}]\frac{(1-x^3)^p}{(1-x)^{p+2}}\\ =&\frac{1}{3^{p-1}}\sum_{0\le k<p/3}\binom{p}{k}(-1)^k\binom{-(p+2)}{p-1-3k}(-1)^{p-1-3k}\\ =&\frac{1}{3^{p-1}}\sum_{0\le k<p/3}\binom{p}{k}(-1)^{p-1-3k}\binom{2p-3k}{p-1-3k}\\ =&\frac{1}{3^{p-1}}\binom{2p}{p-1}+\frac{1}{3^{p-1}}\sum_{0<k<p/3}\binom{p}{k}(-1)^k\binom{2p-3k}{p-1-3k}\\ =&\frac{2p}{3^{p-1}(p-1)}\binom{2p-1}{p-2}+\frac{p}{3^{p-1}}\sum_{0<k<p/3}\frac{(-1)^k}{k}\binom{p-1}{k-1}\binom{2p-3k}{p-1-3k}.\end{aligned}$$

It is known that $\binom{2p-1}{p-2} \equiv (-1)^{p-2} = -1 \pmod{p}$ and

$$\binom{2p-3k}{p-1-3k} = \binom{p+(p-3k)}{p-1-3k} \equiv \binom{p-3k}{p-1-3k} = p-3k \equiv -3k \pmod{p}.$$

Therefore

$$\sum_{k=0}^{p-1}\frac{T_k}{3^k} \equiv 2p+3p\lfloor\frac{p-1}{3}\rfloor = \begin{cases} p \pmod{p^2} & \text{if } p\equiv 1 \pmod 3,\\ 0 \pmod{p^2} & \text{if } p\equiv 2 \pmod 3.\end{cases}$$

We are done. □

Acknowledgment. We are grateful to Professor Zhi-Wei Sun for his very helpful suggestions on our paper.

References

[1] G. E. Andrews, *Euler's "exemplum memorabile inductionis fallacis" and q-trinomial coefficients*, J. Amer. Math. Soc., **3**(1990), 653-669.

[2] G. E. Andrews and R. J. Baxter, *Lattice gas generalization of the hard hexagon model. III. q-trinomial coefficients*, J. Statist. Phys., **47**(1987), 297-330

[3] H. Pan and Z. W. Sun, *A combinatorial identity with application to Catalan numbers*, Discrete Math., **306**(2006), 1921-1940.

[4] Z. W. Sun, *On the sum* $\sum_{k\equiv r (mod m)}\binom{n}{k}$ *and related congruences*, Israel J. Math., **128**(2002), 135-156.

[5] Z. W. Sun and R. Tauraso, *New congruences for central binomial coefficients*, Adv. in Appl. Math., **45**(2010), 125-148.

[6] Z. W. Sun and R. Tauraso, *On some new congruences for binomial coefficients*, Int. J. Number Theory, to appear. http://arxiv.org/abs/0709.1665.

Department of Applied Mathematics, Nanjing Audit University, Nanjing 210029,. People's Public of China

E-mail address: `caohq@nau.edu.cn`

Department of Mathematics, Nanjing University, Nanjing 210093, People's Republic of China

E-mail address: `haopan79@yahoo.com.cn`